arXiv:1303.6853v1 [hep-th] 27 Mar 2013

On the Chern-Simons terms in Lifshitz like quantum electrodynamics

Van Sérgio Alves,[1, *] B. Charneski,[2, †] M. Gomes,[2, ‡]
Leonardo Nascimento,[1, §] and Francisco Peña[3, ¶]

[1]*Faculdade de Física, Universidade Federal do Pará, 66075-110, Belém, Pará, Brazil*
[2]*Instituto de Física, Universidade de São Paulo*
Caixa Postal 66318, 05315-970, São Paulo, SP, Brazil
[3]*Departamento de Ciencias Físicas, Facultad de Ingeniería,*
Ciencias y Administración, Universidad de La Frontera,
Avda. Francisco Salazar 01145, Casilla 54-D, Temuco, Chile

(Dated: March 28, 2013)

Abstract

In this work the generation of generalized Chern-Simons terms in three dimensional quantum electrodynamics with high spatial derivatives is studied. We analyze the self-energy corrections to the gauge field propagator by considering an expansion of the amplitude up to third order in the external momenta. The divergences of the corrections are determined and explicit forms for the Chern-Simons terms with high derivatives are obtained. Some unusual aspects of the calculation are stressed and the existence of a smooth isotropic limit is proved. The transversality of the anisotropic gauge propagator is also discussed.

PACS numbers: 11.10.Gh, 11.10.Hi, 11.10.-z

---

[*] Electronic address: vansergi@ufpa.br
[†] Electronic address: bruno@fma.if.usp.br
[‡] Electronic address: mgomes@fma.if.usp.br
[§] Electronic address: lnascimento@ufpa.br
[¶] Electronic address: fcampos@ufro.cl

I. INTRODUCTION

A great deal of attention has been devoted to the analisis of possible effects of the spacetime anisotropy. These studies involve different areas as gravity and cosmology [1], string theory [2] and condensed matter systems [3]. In particular, in the context of quantum field theory (QFT), this possibility appears as a tool to study nonrenormalizable theories since it improves the ultraviolet behavior of the perturbative series, in spite of violating the Lorentz symmetry. Actually, a lot of Lorentz violating situations has been studied, encompassing noncommutative models [4], the extended standard model [5] and graphene physics [6] and furnishing bounds to Lorentz breaking parameters [7].

The main characteristic of the anisotropic or Lifshitz-like theories is the different behavior between space and time coordinates under a general scaling $x^i \to bx^i$ whereas $t \to b^z t$ [8], where z is the critical exponent; it dictates the renormalizability and the anisotropic behavior of the theory such that a greater value ameliorates the ultraviolet behavior. It was argued that the four-dimensional gravity becomes renormalizable when $z = 3$, for example [9]. However, the implementation of this kind of anisotropy introduces unusual aspects in the general setting of QFT and therefore it is important to carefully investigate the consequences of this new approach [10, 11].

One special situation concerns the Chern-Simons (CS) theories, which have a prominent status on the 2+1 dimensional physics [12]. When the CS term is added to the quantum electrodynamics (QED) Lagrangian, the generation of mass for the gauge field happens without gauge symmetry breaking [12, 13]. Moreover, in gauge theories the CS term naturally emerges from quantum corrections to the gauge field propagator [14]. Beyond that, applications have been devised to condensed matter physics [6, 15] and string theory [16] as well. In particular, in condensed matter physics, the most known effects related to the theories containing the CS term are the quantum Hall effect [17] and superconductivity with high critical temperature [18]. Therefore the study of the influence of the spacetime anisotropy on the theories involving the CS term is certainly relevant for a proper understanding of the 2+1 dimensional physics.

In this work we will analyze the new contributions to the self-energy of the gauge field due to anisotropy in the $z = 2$ case up to one loop order in small momenta regime. In particular, possible corrections to the CS term will be studied. On general grounds we expect that the leading CS corrections have following form:

$$\mathcal{L}_{\mathcal{CS}} \;=\; a\epsilon^{\mu\rho\nu}A_\mu\partial_\rho A_\nu + b\epsilon^{\mu\rho\nu}\Delta A_\mu\partial_\rho A_\nu + c\,\epsilon^{\mu\rho\nu}\partial_0^2 A_\mu\partial_\rho A_\nu, \tag{1}$$

where Δ denotes the Laplacian. In an interesting paper [19], where the isotropic spacetime was considered, a CS term with structure similar to (1) was employed. In that situation the Laplacian was replaced by a D'Alambertian and $c = 0$, so that the gauge propagator shows two massive excitations, one of them being a ghost like.

By starting from (1) with $c = 0$, so that there is at most one time derivative, and adding the Maxwell term,

$$\mathcal{L}_{\mathcal{A}} \;=\; -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{\lambda}{2}(\partial_\mu A^\mu)^2, \tag{2}$$

we obtain the propagator

$$\begin{aligned} D^{\mu\nu}(k) \;=\; & \frac{-i}{[k^2-(a-b\vec{k}^2)^2]}\left(g^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} - \frac{i(a-b\vec{k}^2)\epsilon^{\mu\rho\nu}k_\rho}{k^2}\right) + \\ & - \frac{ik^\mu k^\nu}{\lambda k^2[k^2-(a-b\vec{k}^2)^2]} + \frac{i(a-b\vec{k}^2)^2 k^\mu k^\nu}{\lambda k^2[k^2-(a-b\vec{k}^2)^2]}, \end{aligned} \tag{3}$$

which does not contain particle like poles and, for small momenta, indicates disturbances propagating with squared velocity $(1 - 2ab)$. In this work we will also check the consistency with respect to the gauge symmetry of these corrections through the verification of the transversality of the self-energy correction to the gauge field propagator. Our calculations show that, analogously to the relativistic situation, although finite the coefficient a is regularization dependent.

This work is organized as follows. In Sec. II we introduce high derivative terms in the Dirac and Maxwell Lagrangians and analyze the CS generation and the characteristics of the self-energy of the gauge field. For simplicity, our calculations will be restricted to small momenta regime. In Sec. III we discuss the transversality of the corrections to the gauge field propagator. Sec. IV presents some concluding remarks. Two appendices are dedicated to detail some aspects of the involved calculations.

II. GENERATION OF CHERN-SIMONS TERMS IN THE ANISOTROPIC QED

We begin our study of the effects of the anisotropy on the gauge field propagator and the CS generation by considering the modified Dirac Lagrangian, containing a high spatial derivative of second order

$$\mathcal{L} = \bar{\psi}(i\gamma^0 D_0)\psi + b_1\bar{\psi}(i\gamma^i D_i)\psi + b_2\bar{\psi}(i\gamma^i D_i)^2\psi - M\bar{\psi}\psi, \tag{4}$$

where $i = 1, 2$ and $D^\mu = \partial^\mu - ieA^\mu$ ($\mu = 0, 1, 2$) is the covariant derivative. To obtain the QED Lagrangian we add to (4) the Maxwell term, which contains high derivatives as well

$$\mathcal{L}_{\mathcal{M}} = \frac{1}{4}\left(F_{ij}\Delta F_{ij} + 2F_{0i}F_{0i}\right) = \frac{1}{4}F_{ij}\Delta F_{ij} + \frac{1}{2}\partial_i A_0 \partial_i A_0 - \partial_0 A_i \partial_i A_0 + \frac{1}{2}\partial_0 A_i \partial_0 A_i, \tag{5}$$

in which $F_{ij} = \partial_i A_j - \partial_j A_i$. Note that the above contribution exhibits a mixture among space and time components, which as we mentioned, have distincts scale behavior and may cause complications on the calculations involving the gauge field propagator. It is possible, however, to avoid the mixed propagators by conveniently choosing the gauge fixing [11]. Nevertheless, for our purposes, as the gauge field propagator does not appear in the calculations, that trick is useless and we will keep an inespecific gauge fixing, $\mathcal{L}_{\mathcal{F}}$.

We can rewrite (4) and define the Dirac anisotropic Lagrangian

$$\mathcal{L}_{\mathcal{D}} = \bar{\psi}(i\gamma^0\partial_0)\psi + b_1\bar{\psi}(i\gamma^i\partial_i)\psi + b_2\bar{\psi}(i\gamma^i\partial_i)^2\psi - M\bar{\psi}\psi, \tag{6}$$

where γ^μ indicates a 2×2 representation of the Dirac gamma matrices. Finally, up to irrelevant surface terms, the interaction Lagrangian is

$$\begin{aligned} \mathcal{L}_{\mathcal{I}} &= e\bar{\psi}(\gamma^0 A_0 + b_1\gamma^i A_i)\psi + e^2 b_2\bar{\psi}(\gamma^i A_i)^2\psi + ieb_2(\bar{\psi}\gamma^j\gamma^i\partial_i\psi - \partial_i\bar{\psi}\gamma^i\gamma^j\psi)A_j, \\ &= V_1 + V_2 + V_5 + V_4 + V_3, \end{aligned} \tag{7}$$

such that the total Lagrangian is $\mathcal{L}_{\mathcal{T}} = \mathcal{L}_{\mathcal{M}} + \mathcal{L}_{\mathcal{F}} + \mathcal{L}_{\mathcal{D}} + \mathcal{L}_{\mathcal{I}}$. In (7) we introduced a notation for the vertices where, $V_1 = e\bar{\psi}(\gamma^0 A_0)\psi, V_2 = eb_1\bar{\psi}(\gamma^i A_i)\psi$, etc. These vertices, fixed by $\mathcal{L}_{\mathcal{I}}$, are graphically represented in Fig.(1) in such way that the space and time component are considered separately. Observe that some of the new vertices contain derivatives, wich causes a worsening of the ultraviolet behavior. The anisotropy modifies the kinetic terms of $\mathcal{L}_{\mathcal{T}}$ and consequently the propagators are also changed. For the free fermion propagator we obtain

$$S(k) = \frac{i\left(\hat{\not{k}} + b_1\bar{\not{k}} + b_2\mathbf{k}^2 + M\right)}{k_0^2 - (b_1^2 + 2Mb_2)\mathbf{k}^2 - b_2^2\mathbf{k}^4 - M^2},$$

where the hat and the bar denotes the time and space components, respectively, i.e., $\hat{\not{k}} = \gamma^0 k_0$, $\bar{\not{k}} = \gamma^i k_i$ and $\mathbf{k}^2 = k^i k^i$ (in general we will also use the notations $\hat{k}_0 = k_0$ and $\bar{k}_i = k_i$, thus $\bar{k}^2 = \bar{k}^i\bar{k}_i$).



FIG. 1: Graphical representation of the interaction vertices. The continuos line stands for the fermion field, the dashed and wavy lines are for the A_0 and A_i components, respectively. The diagram (b) may contain a spatial derivative.



FIG. 2: One loop contributions to the gauge self-energy.

To perform the calculation of the corrections to the gauge field propagator at one loop, we consider the graphs represented in Fig.(2) whose analytical expressions are

$$\Pi^{\mu\nu}_{ab} = C_{ab}\int d^z\hat{k}d^d\bar{k}Tr[V^{\mu}_a S(k) V^{\nu}_b S(k+p)], \tag{8}$$

$$\Pi^{ij}_5 = C_5\int d^z\hat{k}d^d\bar{k}Tr[\gamma^i\gamma^j S(k)], \tag{9}$$

where V^{μ}_a is the time or spatial component ($\mu = 0, i$) associated to the vertices defined as in (7), thus $a, b = 1, ..., 4$ and C_{ab} is the momentum independent coefficient correspondent to each amplitude. The development of these expressions are discussed in appendix B.

By power counting we find: the diagram ($2c$) is linearly or logarithmically divergent if the vertex attached to the wavy line has or does not have a derivative; the divergences of the diagram ($2b$) are quadratic if both vertices have one derivative, linear when one vertex has a derivative and none in the other and finally logarithmic when none derivative appears in the vertices just like in the Fig. $(2a)$, which is composed by two external $A^0(p)$ components.

In the sequel we will consider the small momenta regime and Taylor expand the self-energy corrections. The usual CS term (i.e. the first term in (1)) is obtained from the first order terms on the Taylor derivative expansion while the extended CS term (which have a form like the second term in (1)) is given by the third order terms. Considering that the highest divergence is quadratically, the extended CS contributions, which are of third order on Taylor expansion, are of course finite. Differently, for the usual CS terms, the algebra reduces the degree of divergence of some diagrams only to logarithmic so that they requires the introduction of a regularization scheme. Similarly to the relativistic theory, there is an ambiguity on the induced CS term (this effect is shown in appendix B).

For the even terms in the momentum expansion, which do not contribute to the CS terms, the integration on momenta furnishes hypergeometric functions, as indicated in the appendix A. In this case, the divergences are characterized by the Schwinger parameter, x, which receives contributions from the hypergeometric and from the coefficients associated to them (see Eq. A2). To integrate on the x parameter, we will power expand the hypergeometric for small x and $d = 2$, until its exponent becomes nonnegative. Observe that, here the isotropic limit, $b_2 \to 0$, cannot be taken singly because it is inconsistent with the small x expansion. However, although the divergent terms individually exhibit poles for $d = 2$, they are canceled when summed to produce the total self-energy correction, allowing for a smooth isotropic limit.

III. TRANSVERSALITY OF THE GAUGE SELF-ENERGY

In the previous section we discussed the effects of the anisotropy of the spacetime on the induction of CS terms and on the gauge field self-energy divergent terms. However, we said nothing about the transversality of the self-energy corrections. To ensure this fundamental property we can appeal to the conservation of the current which guarantees the transversality of the gauge field propagator. Observe that in (6) the term of higher derivative acting on ψ can be, by integration by parts, rewritten as a term with just first order derivatives. This allows us to use the usual form of the Noether's theorem. However, due to the anisotropy, it is convenient to analyze the space and time components separately

$$J^0 = \frac{\partial \mathcal{L}}{\partial \partial_0 \psi} \delta\psi + \delta\bar{\psi} \frac{\partial \mathcal{L}}{\partial \partial_0 \bar{\psi}}, \tag{10}$$

$$J^k = \frac{\partial \mathcal{L}}{\partial \partial_k \psi} \delta\psi + \delta\bar{\psi} \frac{\partial \mathcal{L}}{\partial \partial_k \bar{\psi}}. \tag{11}$$

The global gauge transformation of the fields are $\delta\psi = i\alpha\psi$, $\delta\bar{\psi} = -i\alpha\bar{\psi}$ and $\delta A^\mu = 0$. After calculating the derivatives in (10), (11) and disregarding the parameter $(-\alpha)$ we get

$$J^0 = \bar{\psi}\gamma^0\psi, \tag{12}$$

$$J^k = b_1\bar{\psi}\gamma^k\psi - ib_2\left[(\partial_i\bar{\psi})\gamma^i\gamma^k\psi - \bar{\psi}\gamma^k\gamma^i(\partial_i\psi)\right] + eb_2\left[\bar{\psi}\gamma^i\gamma^k\psi + \bar{\psi}\gamma^k\gamma^i\psi\right]A_i\ .$$

According to the above result, we can write the $\mathcal{L}_I$ in terms of the conserved current

$$\mathcal{L}_I = \frac{e}{2}\left[A_\mu J^\mu_{(A^\rho=0)} + A_\mu J^\mu\right], \tag{13}$$

where in the first term, the argument of the current (A^ρ) must be taken equal to zero.

Now, in computing of the self-energy contributions notice that it is equal to

$$\langle T \frac{\delta \mathcal{S}_I}{\delta A_\mu} \frac{\delta \mathcal{S}_I}{\delta A_\nu} \rangle \propto \langle J^\mu J^\nu \rangle \tag{14}$$

where $\mathcal{S}_I = \int \mathcal{L}_I$ is the interaction action; (14) must be transversal due to the conservation of the current.

IV. CONCLUDING REMARKS

In this paper we studied the effects of the anisotropy of the spacetime on the generation of the CS term and on the self-energy correction to the gauge field propagator. We proved that this correction is transversal by constructing the conserved Noether's current which interacts with the gauge field.

There are two types of CS terms that may be induced by the radiative corrections, the usual CS term which contains just one derivative and the extended CS term with three derivatives. Whereas the second type of CS term is always finite, the usual one presents results which are divergent but these divergences notabily cancel among themselves when the total contribution is considered. Nevertheless, as it happens in the relativistic situation, the coefficient of the usual CS term is regularization dependent and vanishes if dimensional reduction is adopted. In these amplitudes, it is interesting to notice that the explicit b_2 factor coming from the new vertices is canceled because it is multiplied by amplitudes which diverge as $b_2 \to 0$; consequently, in the isotropic limit there are new contibutions which produce unexpected results (one example is the just mentioned vanishing of the usual CS term if dimensional reduction is employed). On the other hand, as all amplitudes leading to the induced extended CS terms are finite even if $b_2 \to 0$, the isotropic limit for these terms is smooth remaining only the contributions coming from the usual vertices.

By considering the Eqs. (B32) and (B33), the induced CS terms can be written as

$$\Pi^{\mu\nu}_{CS} = -\frac{\left[\alpha_{\mu\nu} + \hat{p}^2 + \bar{p}^2(b_1^2 + 4mb_2)\right] e^2 b_1^2 \epsilon^{\mu\nu\rho} p_\rho}{48\pi m^2 (b_1^2 + 4mb_2)} \tag{15}$$

in accord with our proposal (1). Notice that the breaking of the Lorentz invariance is a very simple function of b_1 and b_2 and that in the isotropic limit with $b_1 = 1$ the Lorenz symmetry is restored.

Acknowledgments

This work was partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenacão de Aperfeiçoamento de Pessoal de Nível Sueprior (CAPES) and Fundacão de Amparo à Pesquisa do Estado de São Paulo (FAPESP). (FP) acknowledge the support to this research by Dirección de Investigación y Desarrollo de la Universidad de La Frontera and Facultad de Ingeniería, Ciencias y Administración, Universidad de La Frontera (Temuco-Chile) .

Appendix A: The calculation procedure

In this appendix we will describe the procedure employed to calculate the Feynman diagrams. As an example, consider the third contribution shown in the appendix B

$$\Pi^{ij}_{22}(p) = \frac{(eb_1)^2}{(2\pi)^3} \int d\hat{k} d^d\bar{k} \; \mathrm{Tr}[\gamma^i S(k) \gamma^j S(k-p)]. \tag{A1}$$

To solve (A1) we adopt the dimensional reduction scheme, in which all the algebra of the gamma matrices are done in $d = 2$ and afterwards the integral is promoted to d dimensions [20]. Therefore, the leading term in the Taylor expansion, i.e. the term with $(\hat{p}=0, \bar{p}=0)$ is

$$-\frac{(eb_1)^2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\left[\frac{2g^{ij}[M^2-\hat{k}^2-b_2\mathbf{k}^2(-\mathbf{k}^2b_2-2M)]}{(\hat{k}^2-\mathbf{k}^2(b_1^2+2Mb_2)-\mathbf{k}^4b_2^2-M^2)^2}\right].$$

To perform the above integral, we use the Schwinger's representation and integrate on the momenta. The result is very extensive, therefore we will consider just one of the terms which leads to a divergent result:

$$e^{-iM^2x}x^{-d/4}\ {}_1F_1\left(\frac{d+2}{4};\frac{1}{2};\frac{ix(b_1^2+2Mb_2)^2}{4b_2^2}\right), \tag{A2}$$

where x is the Schwinger parameter and ${}_1F_1$ denotes the hypergeometric function. Note that this function is divergent in the limit $x\to 0$, so we will Taylor expand it, up to get a nonnegative power of x for $d=2$. Thus we obtain

$$x^{-d/4}\ {}_1F_1\left(\frac{d+2}{4};\frac{1}{2};\frac{ix(b_1^2+2Mb_2)^2}{4b_2^2}\right)\to x^{-d/4}\left(1+2\frac{d+2}{4}\frac{ix(b_1^2+2Mb_2)^2}{4b_2^2}\right) \tag{A3}$$

The final result (B5) is obtained by integrating on x and expanding the result around $d=2$. Observe from the argument of the hypergeometric function in (A2) that the isotropic limit, $(b_2\to 0)$, is incompatible with the adopted expansion for $x\to 0$.

On the other hand, if we consider the subleading term in the Taylor expansion, responsible for the usual CS term, i.e. $\left.\frac{\partial\Pi_{22}^{ij}}{\partial\hat{p}}\right|_{\hat{p}=\bar{p}=0}$, the result is

$$\frac{(eb_1)^2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\left[\frac{2i\epsilon^{ij0}(M-b_2\bar{k}^2)}{(\hat{k}^2-\mathbf{k}^2(b_1^2+2Mb_2)-\mathbf{k}^4b_2^2-M^2)^2}\right]. \tag{A4}$$

To compute the above integral, which is finite by power counting, we simply take $d=2$ and integrate on the momenta.

Appendix B: Interaction vertices

In this appendix we will consider the one loop contributions, for small momenta, to the self-energy of the gauge field classifying each graph by its vertices. In particular, for the usual induced CS terms we analyze its regularization dependence.

1. Graph containing the vertices $(e\bar{\psi}\gamma^0\psi A_0)$ and $(e\bar{\psi}\gamma^0\psi A_0)$

$$\Pi_{11}^{00}=\frac{e^2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\ \mathrm{Tr}[\gamma^0S(k)\gamma^0S(k-p)] \tag{B1}$$

Although this amplitude is logarithmically divergent, after performing the algebra, a total cancellation among the divergent terms occurs furnishing a finite result. Besides that no CS term is induced here.

2. Vertices $(e\bar{\psi}\gamma^0\psi A_0)$ and $(eb_1\bar{\psi}\gamma^i\psi A_i)$

$$\Pi_{12}^{0i} = \frac{e^2 b_1}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\ \mathrm{Tr}[\gamma^0 S(k)\gamma^i S(k-p)]$$

This amplitude also furnishes a finite result and the induced CS term is obtained from one spatial derivative on its Taylor expansion and is given by

$$-\frac{e^2 b_1^2 \epsilon^{0ia}\bar{p}_a}{4\pi(b_1^2+4mb_2)}, \tag{B2}$$

for three space derivatives we obtain

$$-\frac{e^2 b_1^2\,(b_1^4+10m^2b_2^2+6mb_2b_1^2)\,\epsilon^{0ia}\bar{p}_a\bar{p}^2}{48\pi m^2(b_1^2+4mb_2)^2}, \tag{B3}$$

for one space and two time derivatives the result is

$$-\frac{e^2 b_1^2\,(b_1^2+6mb_2)\,\epsilon^{0ia}\bar{p}_a\hat{p}^2}{48\pi m^2(b_1^2+4mb_2)^2}. \tag{B4}$$

3. Vertices $(eb_1\bar{\psi}\gamma^i\psi A_i)$ and $(eb_1\bar{\psi}\gamma^j\psi A_j)$

$$\Pi_{22}^{ij} = \frac{(eb_1)^2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\ \mathrm{Tr}[\gamma^i S(k)\gamma^j S(k-p)]$$

Differently from the previous cases, this amplitude exhibits a divergent result which is expressed as a pole at $d=2$:

$$\frac{ie^2 b_1^2 g^{ij}}{2\pi b_2(d-2)} + \text{finite}. \tag{B5}$$

The induced CS term is obtained from one time derivative and is given by

$$-\frac{e^2 b_1^2 \epsilon^{ij0}\hat{p}_0}{2\pi(b_1^2+4mb_2)}, \tag{B6}$$

for two space and one time derivatives we find

$$-\frac{e^2 b_1^2\,(b_1^4+24m^2b_2^2+8mb_2b_1^2)\,\epsilon^{ij0}\hat{p}_0\bar{p}^2}{48\pi m^2(b_1^2+4mb_2)^2}, \tag{B7}$$

and finally, for three time derivatives:

$$-\frac{e^2 b_1^2\,(b_1^2+8mb_2)\,\epsilon^{ij0}\hat{p}_0\hat{p}^2}{48\pi m^2(b_1^2+4mb_2)^2}. \tag{B8}$$

The following case and the amplitudes discussed in the itens 5, 10 and 11 below are linearly divergent by power counting. However, after the trace calculation, the momenta expansion and symmetric integration the degree of divergence is reduced to logarithmic. In these situations, the usual induced CS terms turn out to be finite but regularization dependent, like it happens in the relativistic model. In order to illustrate this fact we will consider the two divergent terms occurring in the amplitude of the iten 4. $\left(\bar{p}^a \frac{\partial}{\partial \bar{p}^a} \Pi^{0i}_{13}\Big|_{\hat{p}=\bar{p}=0}\right)$:

$$\frac{ie^2 b_2^3}{4\pi^3}\,\epsilon^{0ia}\bar{p}_a \int d^z\hat{k} d^d\bar{k} \frac{b_2^2\bar{k}^8 + 3\hat{k}^2\bar{k}^4}{[\hat{k}^2 + (b_1^2 + 2mb_2)\bar{k}^2 - b_2^2\bar{k}^4 - m^2]^3}. \tag{B9}$$

The above integral was calculated by two different approaches. First we used the Schwinger parametrization,

$$\frac{1}{\left[\hat{k}^2 + (b_1^2 + 2mb_2)\bar{k}^2 - b_2^2\bar{k}^4 - m^2\right]^y} = \frac{i^{-y}}{\Gamma(y)}\int_0^\infty dx\; x^{y-1} e^{ix(\hat{k}^2 + (b_1^2+2mb_2)\bar{k}^2 - b_2^2\bar{k}^4 - m^2)}$$

and the dimensional regularization, which produced the result:

$$\frac{e^2(b_1^2 + 2mb_2)(b_1^2 + 6mb_2)\epsilon^{0ia}\bar{p}_a}{32\pi(b_1^2 + 4mb_2)^2}. \tag{B10}$$

As the second approach, we considered $d = 2$ and integrate on the time component of the momentum. To integrate on the spatial components we employed a cutoff regularization, Λ_1 and Λ_2, to the first and second terms, respectively, in (B9). Thus, we get

$$\left[3\ln(\Lambda_2/\Lambda_1) + \frac{(b_1^4 + 14m^2b_2^2 + 8mb_2b_1^2)}{(b_1^2 + 4mb_2)^2}\right]\frac{e^2\epsilon^{0ia}\bar{p}_a}{16\pi}. \tag{B11}$$

The difference between the two regularization schemes (B10) and (B11) is

$$\left[6\ln(\Lambda_2/\Lambda_1) + 1\right]\frac{e^2\epsilon^{0ia}\bar{p}_a}{32\pi}, \tag{B12}$$

such that this residue can be eliminated by redefining the cutoff ($\Lambda_2 \to e^{-1/6}\Lambda_1$). This is an example of the ambiguity which appears in the calculation of the usual CS term.

4. Vertices $(e\bar{\psi}\gamma^0\psi A_0)$ and $(-ieb_2\partial_i\bar{\psi}\gamma^i\gamma^j\psi A_j)$

$$\Pi^{0i}_{13} \;=\; \frac{e^2 b_2}{(2\pi)^3}\int d\hat{k} d^d\bar{k}\; \mathrm{Tr}[\gamma^0 S(k)\gamma^i\gamma^j S(k-p)(k_i)]$$

This amplitude does not have a pole-like contributions and the induced CS terms are generated by one space derivative:

$$\frac{e^2 b_1^2 \epsilon^{0ia} \bar{p}_a}{8\pi(b_1^2 + 4mb_2)}, \tag{B13}$$

by three space derivatives

$$-\frac{e^2 b_1^2 b_2 \left(b_1^2 + 3mb_2\right) \epsilon^{0ia} \bar{p}_a \bar{p}^2}{48\pi m(b_1^2 + 4mb_2)^2}, \tag{B14}$$

by one space and two time derivatives

$$\frac{e^2 b_1^2 b_2 \hat{p}^2 \epsilon^{0ia} \bar{p}_a}{48\pi m(b_1^2 + 4mb_2)^2}. \tag{B15}$$

Notes that in the computation of (B13) (and also for all the usual CS terms) the explicit b_2 factor coming from the vertex was canceled by the amplitude which becomes divergent when $b_2 \to 0$.

5. Vertices $(e\bar{\psi}\gamma^0\psi A_0)$ and $(ieb_2\bar{\psi}\gamma^j\gamma^i\partial_i\psi A_j)$

$$\Pi_{14}^{0i} = \frac{e^2 b_2}{(2\pi)^3} \int d\hat{k} d^d \bar{k} \ \mathrm{Tr}[\gamma^0 S(k+p)\gamma^j\gamma^i S(k)(k_i)]$$

This amplitude does not exhibit regularization poles and the CS term induced by one space derivative is given by

$$\frac{e^2 b_1^2 \epsilon^{0ia} \bar{p}_a}{8\pi(b_1^2 + 4mb_2)}, \tag{B16}$$

for three space derivatives we have

$$-\frac{e^2 b_1^2 b_2 \left(b_1^2 + 3mb_2\right) \epsilon^{0ia} \bar{p}_a \bar{p}^2}{48\pi m(b_1^2 + 4mb_2)^2}, \tag{B17}$$

for one space and two time derivatives we obtain

$$\frac{e^2 b_1^2 b_2 \hat{p}^2 \epsilon^{0ia} \bar{p}_a}{48\pi m(b_1^2 + 4mb_2)^2}. \tag{B18}$$

6. Vertices $(eb_1\bar{\psi}\gamma^i\psi A_i)$ and $(-ieb_2\partial_m\bar{\psi}\gamma^m\gamma^n\psi A_n)$

$$\Pi_{23}^{ij} = \frac{e^2 b_1 b_2}{(2\pi)^3} \int d\hat{k} d^d \bar{k} \ \mathrm{Tr}[\gamma^i S(k)\gamma^m\gamma^n S(k-p)(k_m)]$$

In this case the pole part is

$$-\frac{ie^2 b_1^2 g^{ij}}{2\pi b_2(d-2)} + \text{finite}. \tag{B19}$$

For one time derivative we have the following induced CS term

$$\frac{e^2 b_1^2 \epsilon^{ij0} \hat{p}_0}{4\pi(b_1^2+4mb_2)}, \tag{B20}$$

for two space and one time derivatives:

$$\frac{e^2 b_1^2 b_2^2 \epsilon^{ij0} \hat{p}_0 \bar{p}^2}{12\pi(b_1^2+4mb_2)^2}, \tag{B21}$$

for three time derivatives

$$\frac{e^2 b_1^2 b_2 \epsilon^{ij0} \hat{p}_0 \hat{p}^2}{24\pi m(b_1^2+4mb_2)^2}. \tag{B22}$$

The previous amplitude as well as the next one are linearly divergent. However, differently from the cases in the itens 5 and 6, after performing all the algebra, the integrals associated to the induced usual CS terms are finite and thus free from ambiguities.

7. Vertices $(eb_1\bar{\psi}\gamma^i\psi A_i)$ and $(ieb_2\bar{\psi}\gamma^n\gamma^m\partial_m\psi A_n)$

$$\Pi_{24}^{ij} = \frac{e^2 b_1 b_2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\ \mathrm{Tr}[\gamma^n\gamma^m S(k)\gamma^i S(k+p)(k_m)]$$

The pole part is given by

$$-\frac{ie^2 b_1^2 g^{ij}}{2\pi b_2(d-2)} + \text{finite}. \tag{B23}$$

To the induced CS terms generated by one time derivative we have

$$\frac{e^2 b_1^2 \epsilon^{ij0} \hat{p}_0}{4\pi(b_1^2+4mb_2)}, \tag{B24}$$

for two space and one time derivatives

$$\frac{e^2 b_1^2 b_2^2 \epsilon^{ij0} \hat{p}_0 \bar{p}^2}{12\pi(b_1^2+4mb_2)^2}, \tag{B25}$$

for three time derivatives

$$\frac{e^2 b_1^2 b_2 \epsilon^{ij0} \hat{p}_0 \hat{p}^2}{24\pi m(b_1^2+4mb_2)^2}. \tag{B26}$$

8. Vertices $(-ieb_2\partial_i\bar{\psi}\gamma^i\gamma^j\psi A_j)$ and $(-ieb_2\partial_m\bar{\psi}\gamma^m\gamma^n\psi A_n)$

$$\Pi_{33}^{ij} = \frac{e^2 b_2^2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\ \mathrm{Tr}[\gamma^i\gamma^j S(k)\gamma^m\gamma^n S(k-p)((k-p)_i)(k_m)]$$

The pole part is

$$\frac{ie^2 b_1^2 g^{ij}}{4\pi b_2(d-2)} + \text{finite}. \tag{B27}$$

In this case no CS term is generated.

9. Vertices $(ieb_2\bar{\psi}\gamma^j\gamma^i\partial_i\psi A_j)$ and $(ieb_2\bar{\psi}\gamma^n\gamma^m\partial_m\psi A_n)$

$$\Pi^{ij}_{44} \;=\; \frac{e^2b_2^2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\;\mathrm{Tr}[\gamma^j\gamma^i S(k)\gamma^n\gamma^m S(k-p)(k_i)((k-p)_m)]$$

The pole part is

$$\frac{ie^2b_1^2g^{ij}}{4\pi b_2(d-2)} + \text{finite} \tag{B28}$$

and no CS term is induced.

10. Vertices $(-ieb_2\partial_i\bar{\psi}\gamma^i\gamma^j\psi A_j)$ and $(ieb_2\bar{\psi}\gamma^m\gamma^n\partial_n\psi A_m)$

$$\Pi^{ij}_{34} \;=\; \frac{e^2b_2^2}{2(2\pi)^3}\int d\hat{k}d^d\bar{k}\;\mathrm{Tr}[\gamma^m\gamma^n S(k)\gamma^i\gamma^j S(k+p)(k_n)(k_i)]$$

To the divergent part we obtain

$$\frac{ie^2b_1^2g^{ij}}{4\pi b_2(d-2)} + \text{finite}. \tag{B29}$$

This amplitude does not generate CS terms.

11. Vertices $(ieb_2\bar{\psi}\gamma^j\gamma^i\partial_i\psi A_j)$ and $(-ieb_2\partial_n\bar{\psi}\gamma^n\gamma^m\psi A_m)$

$$\Pi^{ij}_{43} \;=\; \frac{e^2b_2^2}{2(2\pi)^3}\int d\hat{k}d^d\bar{k}\;\mathrm{Tr}[\gamma^j\gamma^i S(k)\gamma^n\gamma^m S(k-p)(k_n)(k_i)]$$

The pole part is given by

$$= \frac{ie^2b_1^2g^{ij}}{4\pi b_2(d-2)} + \text{finite}. \tag{B30}$$

No CS term is induced in this case.

12. Vertice $(e^2b_2\bar{\psi}\gamma^i\gamma^j\psi A_iA_j)$

$$\Pi^{ij}_{06} \;=\; -\frac{2ie^2b_2}{(2\pi)^3}\int d\hat{k}d^d\bar{k}\;\mathrm{Tr}[\gamma^i\gamma^j S(k)]$$

The pole part resultant from the amplitude is

$$-\frac{ie^2b_1^2}{2\pi b_2(d-2)}\;g^{ij} + \text{finite} \tag{B31}$$

and no CS term is induced.

Adding the corresponding contributions shown in the itens 1 to 12, we find that the divergences are entirely canceled and for the induced CS terms we obtain

$$\Pi^{ij}_{CS} = -\frac{\left[\alpha_{ij} + \bar{p}^2(b_1^2 + 4mb_2) + \hat{p}^2\right] e^2 b_1^2 \epsilon^{ij0}\hat{p}_0}{48\pi m^2(b_1^2 + 4mb_2)} \tag{B32}$$

and

$$\Pi^{0i}_{CS} = -\frac{\left[\alpha_{0i} + \bar{p}^2(b_1^2 + 4mb_2) + \hat{p}^2\right] e^2 b_1^2 \epsilon^{0ia}\bar{p}_a}{48\pi m^2(b_1^2 + 4mb_2)} \tag{B33}$$

where α_{ij} and α_{0i} are constant parameters introduced in (B32) and (B33), respectively, to denote the ambiguity introduced by the regularization scheme. We may note that there is a cancellation of the usual CS term leaving only regularization dependent terms. This occurs due to the contributions of the new vertices introduced by the anisotropy and by the fact that their b_2 vertex factor is eliminated after performing the momenta integrals as it was discussed in the iten 4.

In the isotropic limit $b_2 \to 0$ of (B32) and (B33) all the extended CS terms coming from the new vertices contributions are cancelled and we get

$$\Pi^{ij}_{CS} = -\frac{e^2\left(\alpha_{ij} + b_1^2\bar{p}^2 + \hat{p}^2\right)\epsilon^{ij0}\hat{p}_0}{48\pi m^2}$$

and

$$\Pi^{0i}_{CS} = -\frac{e^2\left(\alpha_{0i} + b_1^2\bar{p}^2 + \hat{p}^2\right)\epsilon^{0ia}\bar{p}_a}{48\pi m^2}.$$

By taken $b_1 = 1$ we obtain an expression similar to the one introduced in [19].

[1] R. A. Konoplya, Phys. Lett. B **679**, 499 (2009); E. Kiritsis, G. Kofinas, Nucl. Phys. B **821**, 467 (2009); G. Bertoldi, B. A. Burrington and A. W. Peet, Phys. Rev. D **82**, 106013 (2010); K. Goldstein, S. Kachru, S. Prakash, S. P. Trivedi, JHEP **1008**, 078 (2010); F. S. Bemfica and M. Gomes, Phys. Rev. D **84**, 084022 (2011); E. Abdalla and A. M. da Silva, Phys. Lett. B **707**, 311 (2012); H. Lu, Y. Pang, C. N. Pope and J. F. Vazquez-Poritz, Phys. Rev. D **86**, 044011 (2012); H. Liao, J. Chen and Y. Wang, Int. J. Mod. Phys. D **21**, 1250045 (2012); O. Obregon and J. A. Preciado, Phys. Rev. D **86**, 063502 (2012).

[2] R. Gregory, S. L. Parameswaran, G. Tasinato and I. Zavala, JHEP **1012**, 047 (2010); A. Donos and J. P. Gauntlett, JHEP **1012**, 002 (2010); K. Balasubramanian and K. Narayan,

JHEP **1008**, 014 (2010); H. Singh, JHEP **1104**, 118 (2011); W. Chemissany and J. Hartong, Class. Quant. Grav. **28**, 195011 (2011); D. Cassani and A. F. Faedo, JHEP **1105**, 013 (2011); K. Narayan, Phys. Rev. D **85**, 106006 (2012); P. Dey and S. Roy, JHEP **1206**, 129 (2012).

[3] D. T. Son, Nucl. Phys. Proc. Suppl. **195**, 217 (2009); S. A. Hartnoll, Class. Quant. Grav. **26**, 224002 (2009); R. G. Cai and H. Q. Zhang, Phys. Rev. D **81**, 066003 (2010); E. J. Brynjolfsson, U. H. Danielsson, L. Thorlacius and T. Zingg, J. Phys. A **43**, 065401 (2010); D. Momeni, M. R. Setare and N. Majd, JHEP **1105**, 118 (2011); S. J. Sin, S. S. Xu and Y. Zhou, Int. J. Mod. Phys. A **26**, 4617 (2011).

[4] M. R. Douglas and N. A. Nekrasov, Rev. Mod. Phys. **73**, 977 (2001); B. Charneski, A. F. Ferrari and M. Gomes, J. Phys. A **40**, 3633 (2007); M. Gomes, V. G. Kupriyanov and A. J. da Silva, Phys. Rev. D **81**, 085024 (2010); M. Gomes, V. G. Kupriyanov and A. J. da Silva, J. Phys. A **43**, 285301 (2010).

[5] D. Colladay and V. A. Kostelecky, Phys. Rev. D **58**, 116002 (1998).

[6] B. Charneski, M. Gomes, T. Mariz, J. R. Nascimento and A. J. da Silva, Phys. Rev. D **79**, 065007 (2009).

[7] V. A. Kostelecky, N. Russell, Rev. Mod. Phys. **83**, 11 (2011); B. Charneski, M. Gomes, R. V. Maluf and A. J. da Silva, Phys. Rev. D **86**, 045003 (2012).

[8] D. Anselmi and M. Halat, Phys. Rev. D **76**, 125011 (2007); D. Anselmi, Ann. Phys. 324, 874 (2009); D. Anselmi Ann. Phys. 324, 1058 (2009); M. Visser, Phys. Rev. D **80**, 025011 (2009).

[9] P. Horava, Phys. Rev. D **79**, 084008 (2009); JHEP **0903**, 020 (2009).

[10] R. Iengo, J. G. Russo and M. Serone, JHEP **0911**, 020 (2009); R. Iengo and M. Serone, Phys. Rev. D **81**, 125005 (2010); P. R. S. Gomes and M. Gomes, Phys. Rev. D **85**, 065010 (2012); P. R. S. Gomes and M. Gomes, Phys. Rev. D **85**, 085018 (2012); K. Farakos and D. Metaxas, Phys. Lett. B **711**, 76 (2012); J. Alexandre, J. Brister and N. Houston, Phys. Rev. D **86**, 025030 (2012).

[11] C. F. Farias, M. Gomes, J. R. Nascimento, A. Yu. Petrov and A. J. da Silva, Phys. Rev. D **85**, 127701 (2012).

[12] S. Deser, R. Jackiw and S. Templeton, Phys. Rev. Lett. **48**, 975 (1982); S. Deser, R. Jackiw and S. Templeton, Ann. Phys. **140**, 372 (1982).

[13] J. S. Schonfeld, Nucl. Phys. **B185**, (1981); R. Jackiw, Phys. Rev. **D29**, 2375 (1984).

[14] A. N. Redlich, Phys. Rev. Lett. **52**, 18 (1984); Phys. Rev. **D29**, 2366 (1984).

[15] A. Foerster and H. O. Girotti, Nucl. Phys. B **342**, 680 (1990); E. Fradkin and A. Lopez, Phys. Rev. B **44**, 5246 (1991).

[16] E. Witten, Prog. Math. **133**, 637 (1995); M. Marino, Rev. Mod. Phys. **77**, 675 (2005); K. Balasubramanian and J. McGreevy, Class. Quant. Grav. **29**, 194007 (2012).

[17] R. E. Prange and S. M. Girvin, *The Quantum Hall Effect*, Springer, Berlin, (1987); T. Chkraborty and P. Pietilainen, *The Fractional Quantum Hall Effect*, Springer Verlag, (1988); E. Fradkin and A. Lopez, Phys. Rev. B **44**, 5246 (1991).

[18] R. B. Laughlin, Phys. Rev. **B23**, 5632 (1981); B. Halperin, Phys. Rev. **B25**, 2185 (1982); F. Wilczek and A. Zee, Phys. Lett. **51**, (1983) 2250; Y. Wu and A. Zee, Phys. Lett. **147B** (1984) 325; F. Wilczek, E. Witten and B. I. Halperin, Int. J. Mod. Phys. **B3**, 1001 (1989); F. Wilczek, *Fractional Statistics and Anyon Superconductivity*, World Scientific, Singapore, (1990); X. G. Wen and A. Zee, Phys. Rev. **B41**, 240 (1990); J. D. Lykken and J. Sonnenshein and N. Weiss, Int. J. Mod. Phys. **A6**, 5155 (1991).

[19] S. Deser and R. Jackiw, Phys. Lett. B **451**, 73 (1999).

[20] W. Siegel, Phys. Lett. B **84**, 193 (1979).